

Psychosocial consequences of widespread of torture and sociopolitical pressure in Iran

Siroos Mirzaei, Hodayoun Alizadeh, Seyed Zarei, Reem Alksiri

Abstract

Background: Violence and political repression in general may have a complex psychosocial impact on societies. Studying these effects in countries with a high grade of repression based on reliable data from inside the country is difficult, mainly due to risks facing researchers. **Methods and results:** The authors developed a strategy to integrate available data from inside Iran, exiled survivors, and reliable international sources to investigate the psychosocial situation in the country, where torture and execution are used frequently. Data on human rights violations, including violence against women, persecution of political activists, and torture and execution of teenagers indicate pervasive strategy of repression in Iran in the last four decades. This thereby has caused substantial long term public mental health impact and continue to cause other social and economic problems and to create a burden on the Iranian population. **Conclusion:** Although, general psychosocial

health of a society is determined by multiple factors, we conclude from the very limited data available from inside Iran that political repression is not the only such factor, but can be assumed to be at least contributing significantly to the increasing physical and mental health problems in the country, creating a difficult position for health professionals especially in detention facilities. **Key words:** Iran, mental health, torture, repression, trauma.

Introduction

Violence and political repression may have a psychosocial impact on different levels of a society. Studying these effects in countries with a high degree of repression such as Iran is difficult from inside the country. A number of international organizations, including the World Medical Association (WMA) and UN institutions have nonetheless provided an objective and frightening picture of the situation in this country.

Iran signed the CCPR - International Covenant on Civil and Political Rights - prior the present regime (4.4.1968) but has after change to the present regime not signed standard humanitarian documents such as the UN Convention against Torture, nor the important OPCAT (Optional Protocol to the Convention against Torture and Other Cruel, Inhuman or Degrading Treatment or Punishment) (1).

The United Nations (UN) has repeatedly voiced concern regarding widespread indiscriminate use of the death penalty in Iran. Reports from the UN Special Rapporteur on the Situation of Human Rights in the Islamic Republic of Iran indicate that prisoners of conscience, including journalists, individuals with dual citizenship, and protesters are being detained arbitrarily on vaguely worded charges, held in poor condition and denied access to health care (2). This represents a violation of the Minimum

Siroos Mirzaei.

Organization for Support of Victims of Torture and War, Viena, Austria.

Correo-e: siroos.mirzaei@wienkav.at

Hodayoun Alizadeh. MA.

Independent HR expert, Viena, Austria.

Correo-e: homal300@gmail.com

Seyed Zarei.

Hemayat Organization for Support of Victims of Torture and War, Viena, Austria.

Correo-e: zarei.seyed33@yahoo.com

Reem Alksiri.

Independent HR expert, Viena, Austria.

Received: July, 2019.

Accepted: November, 2020.

Conflict of interests: none

Statement of Human Rights: This publication does not involve any human participants or animals.

Standard rules for the treatment of prisoners (Mandela rules).

Additionally, the Iranian Government has over decades denied all nominated UN- Special Rapporteurs on Human Rights in Iran access to the country (2).

Obviously, there are nearly no research data available on the prevalence and respective impact of different aspects of persecution in Iran from inside the country. In this manuscript, authors follow a review of secondary independent sources such as UNHCR and Amnesty International, in addition to selected peer reviewed publications related to the discussed issues from inside Iran and from exiled survivors. Available data and observations show that torture and execution are used frequently (2), and a high corruption index (rank 140/180) has been reported by transparency international (<https://www.transparency.org/en/countries/iran>).

Restrictions for physicians in following their ethical guidelines

According World Medical Association (WMA) health professionals have been prevented from treating patients, raising concerns about the veracity of documentation related to the cause of death of patients, and physicians have been forced to support clinically inaccurate documentation to hide human rights violations (3).

Persecution of perpetrators in international courts will be difficult as Iran has not ratified the Rome statute that would permit a court on crimes against humanity in situation where the local government ignores investigation and prosecution of crimes, but Universal Jurisdiction might be an effective instrument to persecute perpetrators in other countries (4).

Ahmadreza Djalali is a physician of disaster medicine who has been wrongfully sentenced to death in Iran. He is an Iranian-born resident of Sweden, who teaches at universities in Italy and Belgium, and whose work involves international scientific collaboration. Since his arrest in 2016, he has been refused medical care despite deteriorating health (5).

HIV has been neglected in Iran probably because of the fact that those at risk for this disease are primarily homosexuals and patients with substance abuse, groups who are discriminated against and are at highest risk for

detention or execution. Two physicians, Kamiar and Arash Alaei, who provided medical care for these prisoners have been arrested (2). Iranian doctors face a difficult choice between threats of persecution and violation of national and/or international medical ethics.

Historical events with impact on public health

A total of 30 studies looking for depression in nurses were investigated. The overall prevalence of depression in nurses in a review of 30 studies was 22%, which was higher than the prevalence of depression among the general population. There is a need for psychologically healthy nurses at hospitals (6).

The relatively high prevalence of many indicators of distress and mental health problems in Iran might be related to a number of social, economic or other factors, indicating persecution or a repressive society.

The effects of the long-lasting war between Iran and Iraq (1980-1988) have been partially studied in regard to post-traumatic stress disorder (PTSD) in veterans and their family members as secondary victims. The authors of that study found decreased cortisol levels in progeny of veterans after rearranging the groups to reflect previous history of PTSD, possibly indicating to a transgenerational effect of trauma (7). On 17 July 1988, Iran notified the Secretary-General of its formal acceptance of resolution 598, which was already proposed by the UN in 1987 (8). Until this time, Ayatollah Khomeini had blocked all attempts to end the war, which Iraq initiated (9). He used opportunity of the “forced war” to eliminate the political opponents by numerous executions and planned to expand the revolution to neighbor countries (10). In 1988 there was a series of execution of political prisoners. Amnesty International recorded the names of > 4,400 disappeared prisoners during this time (11). Execution is also used on homosexual teenagers and child offenders against strong criticism by the UN bodies, even in those with mental disabilities (2).

Widespread of Torture

Data on the widespread use of torture have been published by the UN (2) and data of Iranian refugees as discussed later confirm the wide spread use of torture.

Our own data from examining of survivors of torture from Iran using the “Istanbul protocol” (12) indicate the use of systematic torture methods such as psychological torture, beating the whole body, suspension, falanga, burns with cigarettes, and sexual torture of women and men. This has been confirmed to a major extent in other studies (13-15).

Busch *et al.* reported that in a group of Iranian refugee survivors of torture different methods were used, including burning (38%), electrical torture, and sexual torture (25%) (13). Dehghan recently reported on the high prevalence of sexual torture of Iranian and Kurdish refugees (14). There are also reports of anal torture with hard objects in men (15). The wide-spread historical use of torture in Iran, including the time prior to “revolution”, is even documented in a dedicated torture museum (16). There have been elsewhere legal prosecution of perpetrators and successful reparation for survivors of other nationalities, but in countries as Iran, such due procedure is close to impossible since torture is practiced by state officials based on religious codes (17).

As the assassinations of Iranian Kurdish leaders in Vienna and Berlin (18) have demonstrated, safety is absent for opposition members even in democratic host countries. Morville followed a group of asylum seekers in Danish asylum centers. Forty-three persons, largely torture survivors, aged 20-50y, including Iranians demonstrated to a considerable degree activities of daily living (ADL) impairment secondary to psychological and physical injuries (19). Priebe *et al.* investigated long-term mental sequelae of torture in the country, and explored differences between treatment-seekers and non-treatment-seekers (20). Thirty-four torture victims suffering from enduring mental sequelae and living in Germany were examined. Depression, anxiety, and somatoform disorders were diagnosed with a high degree of comorbidity, with Post Traumatic Stress Disorder (PTSD) being the most frequent diagnosis. Treatment seekers had a higher level of psychopathology, particularly PTSD symptoms of intrusion and increased arousal.

According to the above data and various reports, torture constitutes an epidemic problem in Iranian prisons. But there is *de facto* no way to follow up on these allegations by the justice system. Two medical doctors, Ramin Pourandarjani

and Abdolreza Soudbakhsh, from Kahrizak prison, were murdered after having examined inmates who had been tortured and raped (21).

Members of minorities constitute a significant part of the prison population. Further, minorities suffer from discrimination in the academics, for example, many Baha’i have been reportedly expelled from universities or their applications of admission are rejected (2).

Systematic Violence against Women

In a patriarchal society, with many restrictions on females, economical pressure, physical and/or sexual violence threaten women's health.

According to the Office of the United Nations High Commissioner for Human Rights (OHCHR) Iran has not ratified the Convention on the Elimination of All Forms of Discrimination against Women (EDAW) (1). According to civil law, women are denied equal rights, including in marriage, divorce, child custody, and there is ongoing repression of them in relation to their objection to compulsory veiling. Women who do not wear a hijab in public can be sentenced to flogging, up to two months in prison, and/or be fined (2). Several defenders of women’s rights have been imprisoned (22).

In this context, immediate removal of disparities and use of all necessary means, including legislative, administrative, budgetary, promotional, and judicial measures in Iran have been recommended to accomplish decrease domestic violence (23), also perpetrators should be punished and support for victims accomplished.

A recent study revealed that it is essential to improve healthcare providers’ performance in recognizing and treating domestic violence, paying attention to other related factors, i.e., training and codifying guidelines (24).

In another study, victims of sexual violence had significant differences not only in mental health but also in terms of sexual arousal, sexual orgasm, sexual lubrication, pain and sexual satisfaction (25).

Reportedly, among women who participated in the Shiraz Women's Health Cohort Study, the most prevalent disorder was social dysfunction (78%), followed by somatic symptoms (62%). Mental disorders were 64% most prevalent among married women. In a total group of 265 participants whose husbands were in prison, 65%

had mental disorders. The results of this study showed a considerably higher prevalence of mental disorders among this group of women affected by wide-spread imprisonment, in comparison with the general population. Policy-makers were suggested to pay greater attention to the mental health status of underprivileged Iranian women (26).

Additionally, violence against women was studied in the Northeastern Iran. Some socioeconomic characteristics, such as educational level, occupational status of men, heavy smoking and drug abuse were associated with the occurrence of violence against one's intimate female partner (27). Furthermore, there are vigilantes who attack women and activists for standing against forced hijab laws.

Psychosocial health in the society and among prisoners:

Health-related quality of life (HRQoL) among elderly Iranian individuals is generally low. Policy-makers have recommended HRQoL strategies among the elderly by providing social, economic, and psychological support (28).

Noorbala *et al.*, studied mental health status of Iranians and found an increase in suspected cases of mental disorders among Iranians over age 15 between 1999 to 2015. An interesting finding of this study was that in 1999, rural residents were more at risk of mental disorders, while in 2015, urban residents were more prone to have them. The risk of possible diagnosis of mental disorders increased with age, and was higher in people 65 and above, as well as in widowed, divorced and illiterate individuals (29). The prevalence of common mental disorders in Iran has increased from 21% in 1999 to 32% in 2015. The authors concluded that it is vital for policymakers and health officials to act in order to improve and maintain mental health status of people who are at risk in the country (29).

Given repression in Iran, the prevalence of psychiatric disorders among 180 Kashan prisoners (mean age 32 ± 9 y) was 43%. The most frequent disorders were major depressive disorder (28 %), PTSD (17%) and substance use disorder (17%). History of head trauma in prisoners with psychiatric disorders was 52% (30). About half of all prisoners suffered from them. Treatment of psychiatric

disorder in this group was therefore “essential for prevention of crime”.

In the following we present some indicators of psychological consequences of repression in the Iranian community:

a) Suicide

At both the individual and population levels, the suicide rate has long been understood to correlate with cultural, social, political, and economic forces. In another review the most frequent cause of attempted suicide was family conflict (32%). Other related factors included marital problems (26%), economic constraints (12%), and educational failures (5%). It was concluded that social factors, such as family conflicts and marital problems, had a noticeable role in Iranian suicide research (31).

Thus, suicide seems to be a growing concern in recent years. According to another report, Iran rates third highest among Islamic countries (32). According to statistics, each day >13 people take their lives by suicide in the country; most are aged 15–35. Economic problems, mental illnesses, cultural obligations, political issues and social pressures are the major factors for suicide in the country (32).

b) Children/Adolescents

There are differences in the prevalence of mental disorders among children and adolescents, compared Iran with other countries. Prevalence rates of obsessive-compulsive disorder (ranging from 1% to 11.9% among Iranian children and adolescents) were higher than other regional countries with reported prevalence rates of 0.03% to 2.6%. Social phobia prevalence rates of 8% to 23.5%, and were higher among Iranian children and adolescents than in four non-Iranian communities reporting prevalence rates of 0.08% to 0.9% (33).

Children on death row face additional stress after legal proceedings violating any internationally accepted standards, including fear of their impending execution and the lack of information regarding the date it will take place. The continuous anticipation of imminent death result in additional suffering for alleged child offenders and their families (2).

c) Addiction

The consumption of illegal drugs has increased in Iran in the last decades. In 2017, officials claimed that the number of drug addicts had doubled in the past six years (34). Alcohol, opium and cannabis are the most frequently used illegal drugs, but there are new emerging problems with anabolic steroids, ecstasy and stimulants. Lifetime rates of opiate use is between 1.2 and 8.6% in different parts of the country (35). The phenomenon of dramatic increase of drug addiction is a very possible consequence of shortcomings in social, economic, cultural and political system in the community. The Iranian Ministry of Health and Welfare declared that eight hundred thousand addicts were registered in the country, with an increase of 100% in only 9 years (36). The UN estimates that opium-based drugs are abused by 1,7%-2,8% of the Iranian population, compared with 0,3% of the population in Western Europe (37).

A high percentage of prisoners are drug-addicted, and they even have a significantly higher suicide rate than the already high suicide rate in the general population (42% vs. 35%) (35).

d) Emotional distress in journalists

Another target group under constant political repression is journalists. Iran ranks 173 out of 180 countries on an index of press freedom. Recently, the psychological stressors on wellbeing of Iranian journalists, had 65.8% being in the diaspora, were studied.

The detected stressors included arrest (41,2%), torture (19,3%), assault (10.5%), intimidation (51.4%) and threats to families (43.1%), which were associated with intrusive and arousal PTSD and depressive symptoms. Almost a third of Iranian journalists used barbiturates regularly (38).

Discussion

Medical staff inside and outside of Iran can play an important role in defending human rights of prisoners. This can be achieved by documentation of possible torture sequelae, beyond the treatment of injuries, as it has been described in Istanbul protocol (12). Health professionals in Iran can face serious repercussions. The National societies of physicians must report the threats to all relevant authorities and also to the World Medical Association and Amnesty International.

There are other corporal punishments (cruel and unusual punishment as prohibited by Article 5 of the Universal Declaration of Human Rights, 1948) which are performed on the streets, such as amputation or flogging (39). All intentional corporal injuries must be documented (12) and reports should be sent to the relevant authorities, such as WMA and UN, as Iran does not permit regular monitoring, i.e., by UN Special Rapporteurs.

Conclusions

As a consequence of the difficulties to carry out social research in Iran, there are very limited scientific reports from inside the country. However, available reports and studies indicate an increase in mental health problems, such as suicide and substance abuse rates –even in younger individuals– as well as violence against women. Furthermore, there is credible evidence of widespread use of torture, execution of teenagers, and other substantial human rights problems on the background of non-compliance and failure to sign international conventions.

We conclude that political repression can be an important factor in the increasing physical and mental health problems found in Iran. Nonetheless, there is hope to educate health care professionals and policy makers on the situation there, to bring local and international pressure to bear on the country's leaders, to achieve a more just and safe country.

References:

1. United Nations Treaty Database. OHCHR 2020. URL: https://tbinternet.ohchr.org/_layouts/15/TreatyBodyExternal/Treaty.aspx?CountryID=81&Lang=en.
2. Situation of human rights in the Islamic Republic of Iran. Annual report of the United Nations High Commissioner for Human Rights and reports of the Office of the High Commissioner and the Secretary-General (A/HRC/40/24, Feb. 2019). United Nations- General Assembly 2019. URL: <https://documents-dds-ny.un.org/doc/UNDOC/GEN/G19/028/77/PDF/G1902877.pdf?OpenElement>

3. Iran denies medical care to quell dissent. *Lancet* 2012; 379:1691-2.
4. The States Parties to the Rome Statute. Accessed 2020 Sep 02. URL: https://asp.icc-cpi.int/en_menus/asp/states%20parties/Pages/the%20states%20parties%20to%20the%20rome%20statute.aspx#1 .
5. Burkle FM. Ahmadreza Djalali, MD, PhD is Dying. *Prehosp Disaster Med* 2020;1-2.
6. Saeedi Shahri SS, Ghashghaee A, Behzadifar M, et al. Depression among Iranian nurses: A systematic review and meta-analysis. *Med J Islam Repub Iran* 2017; 31:130.
7. Yahyavi ST, Zarghami M, Naghshvar F, et al. Relationship of cortisol, norepinephrine, and epinephrine levels with war-induced posttraumatic stress disorder in fathers and their offspring. *Rev Bras Psiquiatr* 2015; 37:93-8.
8. Security Council Resolution 598: Iraq-Islamic Republic of Iran. United Nation- Peacemaker 1987. URL: <https://peacemaker.un.org/iraqiran-resolution598>.
9. Pear R. Khomeini accepts 'poison' of ending war with Iraq; U.N. Sending mission. *NY Times* 1988. URL: <https://www.nytimes.com/1988/07/21/us/khomeini-accepts-poison-of-ending-the-war-with-iraq-un-sending-mission.html>.
10. Geschichte des Iran seit 1979. Wikipedia 2020. URL: https://de.wikipedia.org/wiki/Geschichte_des_Iran_seit_1979.
11. 1988 executions of Iranian political prisoners. Wikipedia 2020. URL: https://en.wikipedia.org/wiki/1988_executions_of_Iranian_political_prisoners.
12. Frewer A, Furtmayr H, Krása K, et al. *Istanbul-Protokoll*. Göttingen: V&R unipress, 2015.
13. Busch J, Hansen SH, Hougen HP. Geographical distribution of torture: An epidemiological study of torture reported by asylum applicants examined at the Department of Forensic Medicine, University of Copenhagen. *Torture* 2015; 25:12-21.
14. Dehghan R. The health impact of (sexual) torture amongst Afghan, Iranian and Kurdish refugees: A literature review. *Torture* 2018; 28:77-91.
15. Mirzaei S, Wenzel T. Anal torture with unknown objects and electrical stimuli *Journal of Trauma and Treatment*. 2013; 2:162.
16. Lake E. Grisly truth of Iran's torture prisons revealed in new museum recreating brutal techniques with dummies. *The Sun* 2017. URL: <https://www.thesun.co.uk/news/5062464/iran-torture-prison-museum/> .
17. Mirzaei S, Hardi L, Wenzel T. How to combat torture if perpetrators are supported by a religious "justification". *Torture*. 2011; 21:173-77.
18. German court implicates Iran leaders in '92 killings. *CNN* 1997. URL: <http://edition.cnn.com/WORLD/9704/10/germany.iran/>.
19. Morville AL, Amris K, Eklund M, et al. A longitudinal study of changes in asylum seeker's ability regarding activities of daily living during their stay in the asylum center. *J Immigr Minor Health* 2015; 17:852-59.
20. Priebe S, Esmaili S. Long-term mental sequelae of torture in Iran--who seeks treatment? *J Nerv Ment Dis* 1997; 185:74-77.
21. Death of Ramin Pourandarjani. Wikipedia 2020. URL: https://en.wikipedia.org/wiki/Death_of_Ramin_Pourandarjani.
22. Women's Rights in Iran. *HRW*. 2015. URL: <https://www.hrw.org/news/2015/10/28/women-s-rights-iran>.
23. Kokabisaghi F. Women's right to health in Iran: Domestic implementation of international human rights law. *Int J Health Plann Manage* 2019; 34:501-9.
24. Yousefnia N, Nekuei N, Farajzadegan Z. The relationship between healthcare providers' performance regarding women experiencing domestic violence and their demographic characteristics and attitude towards their management. *J Inj Violence Res* 2018;10.
25. Tadayon M, Hatami-Manesh Z, Sharifi N, et al. The relationship between function and sexual

- satisfaction with sexual violence among women in Ahvaz, Iran. *Electron Physician* 2018;10: 6608-15.
26. Maharlouei N, Hoseinzadeh A, Ghaedsharaf E, et al. The mental health status and associated factors affecting underprivileged Iranian women. *Asian J Psychiatr* 2014; 12:108-12.
 27. Hajian S, Vakilian K, Mirzaii Najm-abadi K, et al. Violence against women by their intimate partners in Shahroud in northeastern region of Iran. *Glob J Health Sci* 2014; 6:117-30.
 28. Tourani S, Behzadifar M, Martini M, et al. Health-related quality of life among healthyelderly Iranians: a systematic review and meta-analysis of the literature. *Health Qual Life Outcomes*. 2018; 16:18.
 29. Noorbala AA, Bagheri Yazdi SA, Faghihzadeh S, et al. Trends of Mental Health Status in Iranian Population Aged 15 and above between 1999 and 2015. *Arch Iran Med* 2017;20(11 Suppl. 1): S2-S6.
 30. Noorbala AA, Bagheri Yazdi SA, Faghihzadeh S, et al. A Survey on Mental Health Status of Adult Population Aged 15 and above in the Province of Mazandaran, Iran. *Arch Iran Med* 2017;20 (Suppl. 1): S83-S86.
 31. Nazarzadeh M, Bidel Z, Ayubi E, et al. Determination of the social related factors of suicide in Iran: a systematic review and meta-analysis. *BMC Public Health* 2013; 13:4.
 32. Suicide in Iran. URL: https://en.wikipedia.org/wiki/Suicide_in_Iran#cite_note-magetemad-4.
 33. Zarafshan H, Mohammadi MR, Salmanian M. Prevalence of Anxiety Disorders among Children and Adolescents in Iran: A Systematic Review. *Iran J Psychiatry* 2015; 10:1-7.
 34. Nikpour G. Drugs and Drug Policy in the Islamic Republic of Iran Brandeis University. Middle East brief 2018:119.
 35. Mohammadkhani P, Khanipour H, Azadmehr H, et al. Trait mindfulness, reasons for living and general symptom severity as predictors of suicide probability in males with substance abuse or dependence. *Iran J Psychiatry* 2015; 10:56-63.
 36. Janmardi M. Drug situation in Iran, with emphasis on content analysis of newspapers. *International Journal of Sociology and Anthropology* 2015; 7:112-17.
 37. Nikpour G. Drugs and Drug Policy in the Islamic Republic of Iran. Middle East Brief 2018;119. URL: https://www.academia.edu/37029091/Drugs_and_Drug_Policy_in_the_Islamic_Republic_of_Iran.
 38. Feinstein A, Feinstein S, Behari M, et al. The psychological wellbeing of Iranian journalists: a descriptive study. *JRSM Open* 2016; 7:2054270416675560.
 39. Iran: Authorities amputate a man's hand in shocking act of cruelty. *amnesty international* 2018. URL: <https://www.amnesty.org/en/latest/news/2018/01/iran-authorities-amputate-a-mans-hand-in-shocking-act-of-cruelty/>.

